

Arithmetic properties of the adjacency matrix of quadriculated disks

Nicolau C. Saldanha and Carlos Tomei

February 1, 2008

Abstract

Let Δ be a bicolored quadriculated disk with black-to-white matrix B_Δ . We show how to factor $B_\Delta = L\tilde{D}U$, where L and U are lower and upper triangular matrices, $\tilde{D}$ is obtained from a larger identity matrix by removing rows and columns and all entries of L , $\tilde{D}$ and U are equal to 0, 1 or -1 .

1 Introduction

In this paper, a *square* is a topological disk with four privileged boundary points, the *vertices*. A *quadriculated disk* is a closed topological disk formed by the juxtaposition along sides of finitely many squares such that interior vertices belong to precisely four squares. A simple example of a quadriculated disk is the $n \times m$ rectangle divided into unit squares, another example is given in figure 1. A quadriculated disk may be considered as a closed subset of the plane $\mathbb{R}^2$.




Figure 1: A quadriculated disk and its black-to-white matrix

1991 *Mathematics Subject Classification*. Primary 05B45, Secondary 05A15, 05C50, 05E05.
Keywords and phrases Quadriculated disk, tilings by dominoes, dimers.
The authors gratefully acknowledge the support of CNPq, Faperj and Finep.

Quadriculated disks are *bicolored*, i.e., the squares are black and white in a way that squares with a common side have opposite colors. Label the black (resp. white) squares of a quadriculated disk Δ by $1, 2, \dots, b$ (resp. $1, 2, \dots, w$). The $b \times w$ *black-to-white* matrix B_Δ has (i, j) entry $b_{ij} = 1$ if the i -th black and j -th white squares share an side and $b_{ij} = 0$ otherwise. If the disk has a single square of either color the matrix B collapses. See figure 1 for an example of black-to-white matrix; black squares are numbered. Throughout the paper, blank matrix entries equal 0. For a given labeling of squares of a quadriculated disk Δ in which black squares come first, the adjacency matrix is

$$\begin{pmatrix} 0 & B \\ B^T & 0 \end{pmatrix}.$$

A rectangular matrix $\tilde{D}$ is a *defective identity* if it can be obtained from the identity matrix by adding rows and columns of zeroes or, equivalently, by removing rows and columns from a larger identity matrix. More precisely, $\tilde{D}$ is a defective identity if every entry equals 0 or 1 and if $\tilde{d}_{ij} = \tilde{d}_{i'j'} = 1$ implies either $i = i', j = j'$, or $i < i', j < j'$, or $i > i', j > j'$. For a $n \times m$ matrix A , an $L\tilde{D}U$ decomposition of A is a factorization $A = L\tilde{D}U$ where L (resp. U) is $n \times n$ (resp. $m \times m$) lower (resp. upper) invertible and $\tilde{D}$ is a defective identity. The main result of this paper is the following.

Theorem 1 *Let Δ be a quadriculated disk with at least two squares. For an appropriate labeling of its squares, the black-to-white matrix B_Δ admits an $L\tilde{D}U$ decomposition whose factors have all entries equal to 0, 1 or -1 .*

Thus, for example, the matrix B_Δ in figure 1 admits the decomposition

$$\begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 1 & & & \\ 1 & & & -1 & & \\ & & & & 1 & \\ & 1 & & & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 1 & & & \\ & & & 1 & & 0 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & & 1 & & \\ & 1 & 1 & & 1 & 1 \\ & & 1 & & 1 & \\ & & & 1 & & -1 \\ & & & & 1 & 1 \\ & & & & & 1 \\ & & & & & & 1 \end{pmatrix}.$$

The determinant $\det(B_\Delta)$ is, up to sign, the q -counting for $q = -1$ of the domino tilings of Δ , in the sense of [2]. A *domino tiling* of Δ is a decomposition of Δ as a disjoint union of 2×1 rectangles: $b = w$ is a necessary but not sufficient condition for the existence of a domino tiling. The result below was originally proved in [1] using a very different argument.

Corollary 1.1 *Let Δ be a quadriculated disk with $b = w$ and black-to-white matrix B_Δ . Then $\det(B_\Delta)$ equals 0, 1 or -1 .*

Theorem 1 is stronger, however, especially when B_Δ is not invertible.

Corollary 1.2 *Let Δ be a quadriculated disk with black-to-white matrix B_Δ . If v has integer entries and the system $B_\Delta x = v$ admits a rational solution then the system admits an integer solution.*

This corollary may be interpreted as saying that the cokernel $\mathbb{Z}^b/B_\Delta(\mathbb{Z}^w)$ of $B_\Delta : \mathbb{Z}^w \rightarrow \mathbb{Z}^b$ is a free abelian group. From Theorem 1, the rank r of B_Δ is the same in $\mathbb{Q}$ as in $\mathbb{Z}_p$ for any prime number p . The proof of Corollary 1.1 in [1] is based on this fact for $p = 2$.

The proof of Theorem 1 is constructive in the sense that it yields a fast algorithm to obtain the appropriate labeling of vertices, the matrices in the factorization, $\det(B_\Delta)$ and $\text{rank}(B_\Delta)$.

In section 2 we introduce *cut and paste*, a geometric procedure that converts a quadriculated disk Δ into a union of smaller quadriculated disks $\Delta'_1, \dots, \Delta'_\ell$. In section 3 we present its algebraic counterpart, relating the black-to-white matrices $B_{\Delta'_1}, \dots, B_{\Delta'_\ell}$ to the original B_Δ ; this will be the inductive step in the proof of Theorem 1. In section 4 we study *boards*, quadriculated disks which are subsets of the quadriculated plane $\mathbb{Z}^2 \subset \mathbb{R}^2$, and show that cut and paste can be performed within this smaller class (Theorem 2).

We would like to thank Tania V. de Vasconcelos for helpful conversations and the referee for a careful report which led to a clearer and more correct paper.

2 Cut and paste

The proof of Theorem 1 makes use of *good diagonals*: examples are given in figure 2. *Cut and paste* along a good diagonal obtains from a quadriculated disk Δ a disjoint union of smaller disks $\Delta'_1, \dots, \Delta'_\ell$, often with $\ell = 1$ (Lemma 2.3). Our argument consists of two steps: a proof that there always exists a good diagonal (Proposition 2.2) and a procedure to convert $L\tilde{D}U$ decompositions of $B_{\Delta'_1}, \dots, B_{\Delta'_\ell}$ into a decomposition of B_Δ (Lemma 3.2).

Topological subdisks of $\mathbb{R}^2$ consisting of unit squares with vertices in $\mathbb{Z}^2$ are *boards*: the quadriculated disk in figure 1 is not a board. Arbitrary quadriculated disks can still be “ironed” so that its squares become unit squares with vertices in $\mathbb{Z}^2$: this is what *developing maps* make precise (see [4] for a more general discussion of developing maps).

Lemma 2.1 *Let Δ be a quadriculated disk: there exists a continuous map $\phi : \Delta \rightarrow \mathbb{R}^2$ which is locally injective in the interior of Δ and takes squares in Δ to unit squares in $\mathbb{R}^2$ with vertices in $\mathbb{Z}^2$.*

Proof: Orient Δ and endow it with a metric structure such that each square becomes isometric to the unit square in the plane: ϕ is to be an orientation preserving local isometry in the interior of Δ . Pick an arbitrary edge v_0v_1 and set $\phi(v_0) = (0, 0)$, $\phi(v_1) = (1, 0)$ and extend ϕ isometrically along the edge. Notice that an orientation preserving local isometry extends uniquely from an edge to the squares adjacent to the edge. By extending the domain of ϕ repeatedly starting from v_0v_1 , uniqueness is clear. Since Δ is simply connected, ϕ is well-defined. ■

For the quadriculated disk in figure 1, the developing map ϕ is neither injective in the interior of Δ nor locally injective on its boundary.

There exist genuinely different quadriculated disks Δ and $\tilde{\Delta}$ for which the image of the restriction of the developing map ϕ to their boundaries is the same: this a corollary of Milnor's paisley curve example ([3]). It is not hard to produce such examples with $\det(B_\Delta) = 0 \neq \det(B_{\tilde{\Delta}})$.

A *corner* of a quadriculated disk is a boundary point which is a vertex of a single square. A *diagonal* of a quadriculated disk is a sequence of vertices $v_0v_1 \dots v_k$, $k > 0$, such that (i) v_0 is a corner, $v_1, v_2, \dots, v_{k-1}$ are interior vertices and v_k is a boundary vertex; (ii) consecutive vertices v_i and v_{i+1} , $i = 0, \dots, k-1$, are opposite vertices of a square s_{i+1} ; (iii) consecutive squares s_i and s_{i+1} , $i = 1, \dots, k-1$, have a single vertex in common (which is v_i).

Notice that both coordinates of $\phi(v_i)$ depend monotonically on i . In particular, vertices and squares of a diagonal of a quadriculated disk are all distinct. A diagonal $v_0 \dots v_k$ is a *good diagonal* if at least one side of the square s_k containing v_k is contained in the boundary of Δ . A good diagonal $v_0 \dots v_k$ is *balanced* (see figure 2) if exactly one side of s_k containing v_k is contained in the boundary of Δ , and *unbalanced* otherwise.




Figure 2: Good balanced, bad and good unbalanced diagonals

Proposition 2.2 *Any quadriculated disk Δ admits at least four good diagonals.*

Notice that diagonals, being sequences of vertices, are naturally oriented. In particular, a square has four diagonals, all good.




Figure 5: Notation for cut and paste

6

Proof: Notice that some isolated points in ζ^r of the form v_i^r may be boundary points but otherwise ζ^r is in the interior of Δ ; the zig-zag ζ^{l+} , on the other hand, may contain boundary segments (see figure 6). To prove path connectivity of Δ^r , take the shortest path in Δ connecting an arbitrary point in Δ^r to a point in ζ^r . If a simple closed curve is contained in Δ^r (or Δ^l) then so is the disk surrounded by the curve: this prove simple connectivity of the connected components. Since Δ^l , Δ^r and their intersection $\zeta^l = \zeta^r$ in $\tilde{\Delta}'$ are all path connected and simply connected, so is $\tilde{\Delta}'$. Interior vertices of $\tilde{\Delta}'$ are surrounded by four squares and therefore $\tilde{\Delta}'$ is quadriculated. ■




Figure 6: Cut and paste may produce a disjoint union of disks

With the notation of the lemma, set Δ' to be the disjoint union of the quadriculated disks $\Delta'_1, \dots, \Delta'_\ell$. In this sense, cut and paste obtains from a disk a (possibly empty) disjoint union of quadriculated disks. Under the obvious extension of the concept of black-to-white matrix,

$$B_{\tilde{\Delta}'} = B_{\Delta'} = \begin{pmatrix} B_{\Delta'_1} & \dots & 0 \\ \vdots & & \vdots \\ 0 & \dots & B_{\Delta'_\ell} \end{pmatrix}$$

for the same labeling of squares and thus $L\tilde{D}U$ decompositions of $B_{\Delta'_1}, \dots, B_{\Delta'_\ell}$ yields a similar decomposition of $B_{\Delta'}$. Figure 6 shows an example of Δ , $\tilde{\Delta}'$ and Δ' where $\ell = 2$.

3 The decomposition

Before discussing the relationship between B_Δ and $B_{\Delta'}$ we present a lemma in linear algebra. The proof is a straightforward computation left to the reader. Let I_n be the $n \times n$ identity matrix and $I_{n,m}$ be the $n \times m$ defective identity matrix with (i, j) entry equal to 1 if $i = j$ and 0 otherwise.

Lemma 3.1 *Let M be a $(n + m) \times (n' + m')$ matrix which can be decomposed in blocks as*

$$M = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix},$$

where M_{11} is $n \times n'$. If $n' \leq n$ and N is a $n' \times m'$ matrix with $M_{11}N = M_{12}$ then

$$M = \begin{pmatrix} M_{11}I_{n',n} & 0 \\ M_{21}I_{n',n} & I_m \end{pmatrix} \begin{pmatrix} I_{n,n'} & 0 \\ 0 & M_{22} - M_{21}N \end{pmatrix} \begin{pmatrix} I_{n'} & N \\ 0 & I_{m'} \end{pmatrix}.$$

Similarly, if $n' \geq n$ and N is a $m \times n$ matrix with $NM_{11} = M_{21}$ then

$$M = \begin{pmatrix} I_n & 0 \\ N & I_m \end{pmatrix} \begin{pmatrix} I_{n,n'} & 0 \\ 0 & M_{22} - NM_{12} \end{pmatrix} \begin{pmatrix} I_{n',n}M_{11} & I_{n',n}M_{12} \\ 0 & I_{m'} \end{pmatrix}.$$

The next lemma is the inductive step in the proof of Theorem 1.

Lemma 3.2 *Let Δ be a quadriculated disk with b black and w white squares, $b+w > 1$. Let $\Delta' = \Delta'_1 \cup \dots \cup \Delta'_\ell$ (with $b' = b'_1 + \dots + b'_\ell$ black and $w' = w'_1 + \dots + w'_\ell$ white squares) be obtained from Δ by cut and paste along a good diagonal. Label black and white squares in Δ so that excised squares come first and in the order prescribed by the good diagonal; label squares in Δ' next. Then the black-to-white matrices B_Δ and $B_{\Delta'}$ satisfy*

$$B_\Delta = \begin{pmatrix} L & 0 \\ X & S_{b'} \end{pmatrix} \begin{pmatrix} I_{b-b', w-w'} & 0 \\ 0 & B_{\Delta'} \end{pmatrix} \begin{pmatrix} U & Y \\ 0 & S_{w'} \end{pmatrix}$$

where L (resp U) is an invertible lower (resp. upper) square matrix of order $b-b'$ (resp. $w-w'$) and $S_{b'}$ and $S_{w'}$ are square diagonal matrices. Furthermore, all entries of $S_{b'}$, $S_{w'}$, L , U , X and Y equal 0, 1 or -1 .

The statement above requires clarification in some degenerate cases. If Δ' is empty, $B_{\Delta'}$ collapses and $B_\Delta = LI_{b,w}U$. If instead Δ' is a disjoint union of unit squares, all of the same color, then either $w' = 0$ or $b' = 0$ and

$$B_\Delta = \begin{pmatrix} L & 0 \\ X & S_{b'} \end{pmatrix} \begin{pmatrix} I_{b-b', w} \\ 0 \end{pmatrix} U \quad \text{or} \quad B_\Delta = L \begin{pmatrix} I_{b, w-w'} & 0 \end{pmatrix} \begin{pmatrix} U & Y \\ 0 & S_{w'} \end{pmatrix}.$$

Proof: Assume that the squares on the good diagonal are black; thus $k = b-b'$; if the squares were white all computations would be transposed. Let $j_1, \dots, j_{k-1}$ be

the indices of the white squares $s_1^r, \dots, s_{k-1}^r$; notice that $j_i > w - w'$. Decompose the matrix B_Δ in four blocks,

$$B_\Delta = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix},$$

where B_{22} is a $b' \times w'$ matrix. By construction, B_{11} has one of the two forms below, the first case corresponding to balanced good diagonals (i.e., to $b - b' = w - w'$).

$$B_{11} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & 1 & 1 \end{pmatrix}, \quad \text{or} \quad B_{11} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & 1 & 0 & \cdots & 0 \\ 0 & 1 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

Let S_b (resp. S_w) be a $b \times b$ (resp. $w \times w$) diagonal matrix with diagonal entries equal to 1 or -1 ; the i -th entry of S_b (resp. S_w) is -1 if the i -th black (resp. white) square is strictly to the right of the diagonal. Write

$$S_b = \begin{pmatrix} I_{b-b'} & 0 \\ 0 & S_{b'} \end{pmatrix}, \quad S_w = \begin{pmatrix} I_{w-w'} & 0 \\ 0 & S_{w'} \end{pmatrix}.$$

We have

$$S_b B_\Delta S_w = \begin{pmatrix} B_{11} & -B_{12} \\ B_{21} & B_{22} \end{pmatrix}.$$

The matrix B_{12} has $k - 1$ nonzero columns in positions $j_1, \dots, j_{k-1}$. More precisely, the nonzero entries are (i, j_i) and $(i + 1, j_i)$ for $i = 1, \dots, k - 1$. The nonzero columns of B_{12} are equal to the first $k - 1$ columns of B_{11} . Let N be the $(w - w') \times w'$ matrix with entries 0 or -1 , with nonzero entries at $(1, j_1), (2, j_2), \dots, (k - 1, j_{k-1})$. Clearly $B_{11}N = -B_{12}$ and we may apply Lemma 3.1 to write $S_b B_\Delta S_w$ as

$$\begin{pmatrix} B_{11}I_{w-w', b-b'} & 0 \\ B_{21}I_{w-w', b-b'} & I_{b'} \end{pmatrix} \begin{pmatrix} I_{b-b', w-w'} & 0 \\ 0 & B_{22} - B_{21}N \end{pmatrix} \begin{pmatrix} I_{w-w'} & N \\ 0 & I_{w'} \end{pmatrix}.$$

We claim that $B_{\Delta'} = B_{22} - B_{21}N$. The nonzero columns of the matrix $-B_{21}N$ are the columns of B_{21} , except that the first column is moved to position j_1 , the second column is moved to j_2 and so on. The k -th column of B_{21} , if it exists, is discarded. These nonzero entries correspond precisely to the identifications which must be performed in order to obtain Δ' , i.e., to the ones which must be added to B_{22} in order to obtain $B_{\Delta'}$. We now have

$$B_\Delta = \begin{pmatrix} B_{11}I_{w-w', b-b'} & 0 \\ S_{b'}B_{21}I_{w-w', b-b'} & S_{b'} \end{pmatrix} \begin{pmatrix} I_{b-b', w-w'} & 0 \\ 0 & B_{\Delta'} \end{pmatrix} \begin{pmatrix} I_{w-w'} & NS_{w'} \\ 0 & S_{w'} \end{pmatrix}.$$

If the good diagonal is balanced, this finishes the proof. In the unbalanced case, $\tilde{L} = B_{11}I_{w-w', b-b'}$ is not invertible since its last column is zero. Replace the (k, k) entry of $\tilde{L}$ by 1 to obtain a new matrix L : L is clearly invertible and $\tilde{L}I_{b-b', w-w'} = LI_{b-b', w-w'}$. The proof is now complete. $\blacksquare$

Proof of Theorem 1: The basis of the induction on the number of squares of Δ consists of checking that the theorem holds for disks with at most two squares. Notice that if the disk consists of a single square then $b = 0$ or $w = 0$ and the matrices are degenerate.

Let Δ be a quadriculated disk and $\Delta' = \Delta'_1 \cup \dots \cup \Delta'_\ell$ be obtained from Δ by cut and paste. By induction on the number of squares the theorem may be assumed to hold for each Δ'_k and we therefore write $B_{\Delta'} = L_{\Delta'} \tilde{D}_{\Delta'} U_{\Delta'}$. From the induction step, Lemma 3.2, write

$$\begin{aligned} B_{\Delta} &= \begin{pmatrix} L_{\text{step}} & 0 \\ X_{\text{step}} & S_{b'} \end{pmatrix} \begin{pmatrix} I_{b-b', w-w'} & 0 \\ 0 & B_{\Delta'} \end{pmatrix} \begin{pmatrix} U_{\text{step}} & Y_{\text{step}} \\ 0 & S_{w'} \end{pmatrix} \\ &= L_{\Delta} \tilde{D}_{\Delta} U_{\Delta}. \end{aligned}$$

where

$$L_{\Delta} = \begin{pmatrix} L_{\text{step}} & 0 \\ X_{\text{step}} & S_{b'} \end{pmatrix} \begin{pmatrix} I_{b-b'} & 0 \\ 0 & L_{\Delta'} \end{pmatrix}, \quad U_{\Delta} = \begin{pmatrix} I_{w-w'} & 0 \\ 0 & U_{\Delta'} \end{pmatrix} \begin{pmatrix} U_{\text{step}} & Y_{\text{step}} \\ 0 & S_{w'} \end{pmatrix}.$$

The theorem now follows from observing that each nonzero entry of L_{Δ} (resp. U_{Δ}) is, up to sign, copied from either $L_{\Delta'}$, L_{step} or X_{step} (resp. $U_{\Delta'}$, U_{step} or Y_{step}) and is therefore equal to 1 or -1 . $\blacksquare$

The example in Figure 7 is instructive: the B_G matrix of this planar graph G has determinant 1 but admits no $L\tilde{D}U$ decomposition where the matrices have integer coefficients since the removal of any two vertices of opposite colors from G yields a graph whose determinant has absolute value greater than 1.




Figure 7: Determinant 1 does not imply $L\tilde{D}U$ decomposition

4 Boards

A *board* is a quadriculated disk Δ for which the developing map $\phi : \Delta \rightarrow \mathbb{R}^2$ is injective. In other words, a board is a topological subdisk of $\mathbb{R}^2$ whose boundary is a polygonal curve consisting of segments of length 1 joining points in $\mathbb{Z}^2$. In this section we always assume ϕ to be the inclusion. The class of boards is not closed under cut and paste: in figure 8, the two enhanced segments on the boundary would be superimposed by cut and paste with the good diagonal on the left. Cut and paste along the good diagonal indicated on the right, however, yields a smaller board.




Figure 8: A board and two good diagonals, one excellent.

Theorem 2 *It is always possible to cut and paste a given board Δ to obtain a disjoint union of boards Δ' .*

Proof: A diagonal is *excellent* if the x and y coordinates are both monotonic along one of the two boundary arcs between v_0 and v_k ; without loss, let this arc lie to the right of the diagonal. Excellent diagonals are good: the vertex $v^r + k - 1/2$ is on the boundary. We interpret cut and paste along an excellent diagonal as leaving Δ^l fixed and moving Δ^r . In this way, Δ' becomes a subset of Δ and is therefore a disjoint union of boards. We are left with proving that any board admits excellent diagonals.

Each diagonal defines two boundary arcs: order these arcs by inclusion. We claim that a diagonal defining a minimal arc is excellent. Indeed, let $\delta^m = (v_0^m v_1^m \dots v_k^m)$ be a diagonal inducing a minimal arc α : assume without loss of generality that $v_i^m = (a+i, b+i)$ for given a and b . Assume by contradiction that δ^m is not excellent. Consider the value of $x+y$ on α : this function is not strictly increasing and therefore admits a local minimum somewhere in the interior of α . The local minimum closest to v_0^m is the starting point for a diagonal defining a boundary arc β strictly contained in α , a contradiction. ■

References

- [1] DEIFT, P. A. AND TOMEI, C., *On the determinant of the adjacency matrix for a planar sublattice*, J. Combin. Theory Ser. B, **35**, 278–289 (1983).
- [2] ELKIES, N., KUPERBERG, G., LARSEN, M. AND PROPP, J., *Alternating sign matrices and domino tilings*, J. Alg. Combin. **1**, 112–132 and 219–239 (1992).
- [3] POÉNARU, V., *Extension des immersions en codimension 1 (d’après Blank)*, Séminaire Bourbaki 1967/1968, Exposé 342 (Benjamin, New York, 1969).
- [4] THURSTON, W. P. AND LEVY, S. (EDITOR), *Three dimensional geometry and topology*, Princeton Univ. Pr., 1997.

Nicolau C. Saldanha and Carlos Tomei
Departamento de Matemática, PUC-Rio
R. Marquês de S. Vicente 225, Rio de Janeiro, RJ 22453-900, Brazil
nicolau@mat.puc-rio.br; <http://www.mat.puc-rio.br/~nicolau/>
tomei@mat.puc-rio.br